



Review Article

The importance of surface tension in endodontic irrigation: A review study

Hamidreza Abdollahi Mansoorkhani^{1*}, Farshid Mahmoudi¹¹Dept. of Endodontics, School of Dentistry, Islamic Azad University of Shiraz, Iran

ARTICLE INFO

Article history:

Received 02-10-2022

Accepted 28-12-2022

Available online 20-02-2023

Keywords:

Surface Tension

Sodium Hypochlorite

Chlorhexidine

Root Canal

ABSTRACT

The surface tension of irrigants determines their ability to penetrate dentinal tubules and access the apical third of root canals. It is defined as an affinity for a liquid's surface area to decline in the case of surface tension between molecules. Irrigating the canal wall is vital for dissolving organic tissues and disinfecting dentin and its tubules in root canal preparation. Even with super elastic rotary preparation, untouched spaces are a typical issue during root canal instrumentation. As a result, the antimicrobial activity and solvent action in these areas is essential to the disinfection of root canals. The ability to diagnose irrigants quickly and treat them successfully depends on a thorough discernment of their surface tension. In this study, surface tension involved in endodontic solutions was examined for etiological factors, features, and possible consequences. The relevant studies were found by searching online databases. We identified 30 articles using the hand search method and summarized the critical factors of each study. It was found that low surface tension agents would penetrate tubules more acceptable and that surface tensions estimated at various temperatures and concentrations affected canal cleaning.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial 4.0 International](https://creativecommons.org/licenses/by-nc/4.0/), which allows others to remix, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Root canal preparation is mainly concerned with cleaning the canal. Treatment failure may be caused by residue pulp tissue, infected dentine, or bacteria within the canal system. It is crucial for endodontic treatment success to use irrigating solutions for root canal space debridement. During the process of enlarging and shaping the root canal system, endodontic irrigants play a pivotal role in cleaning the root canal system. Preparation of the root canal system may be meant to gain radicular access to the un-instrumented root canal system, thereby dispersing irrigants.¹ When a chemical solution is used for endodontic treatment, it must possess substantial wetness, allowing it to expand its solvent capacity and improve its antimicrobial properties.^{2,3} Sodium hypochlorite (NaOCl),

chlorhexidine (CHX), and Biopure MTAD (a mixture of Doxycycline, citric acid, and detergent) are regarded as promising irrigants for root canal irrigation.^{4,5} Endodontic treatments have long used sodium hypochlorite (NaOCl) for this purpose. Aside from being an ideal organic solvent and an effective antimicrobial, it also irritates the periapical tissues, particularly at high concentrations.^{6,7} The biocompatibility of CHX as well as its use in allergy-related bleaching solutions makes it useful for treating open apices, root resorptions, foramen enlargements, and root perforations.⁸⁻¹⁰ Aside from contact dermatitis, CHX can produce discoloration of the teeth and tongue, and in rare cases, dysgeusia. The antibacterial properties of MTAD in endodontic irrigants enable them to remove smear layers. As compared to other irrigant solutions, MTAD is more biocompatible and less cytotoxic.¹¹⁻¹³ A common irrigant used in root canal treatment is ethylenediaminetetraacetic acid (EDTA). Due to its heightened dynamic viscosit,

* Corresponding author.

E-mail address: h.r.abdollahi65@gmail.com (H. A. Mansoorkhani).

Ethylenediaminetetraacetic acid (EDTA) is ineffective in dissolving the smear layer in the apical third of root canals. In recent years, silver nanoparticles (AgNPs) have gained attention in dentistry as root canal irrigants or medicaments.^{14–17} In addition to being highly effective against bacteria, these particles are compatible with human cells^{18–20} Thus, it is vital to investigate the properties of these newly introduced agents in comparison to typical endodontic irrigants.²¹ Identifying factors that contribute to the etiology, management methods, and outcomes in endodontic treatment with dental irrigants surface tension was the purpose of this study.

2. Discussion

2.1. NaOCl

Its bactericidal and lubricant properties, as well as its ability to dissolve organic substances, make NaOCl an ideal endodontic irrigant.²² A 5% NaOCl solution produces a higher surface tension than a 2.5% solution, but it is not superior to the 2.5% solution in terms of cleansing capability; lower concentrations are suggested due to the material's toxicity. A significant difference between exposure times was observed in the depth of NaOCl penetration as the exposure time increased.^{23,24} Increases in hypochlorite concentration typically result in a greater penetration of regular solutions. Even so, dissimilarities were not statistically noteworthy in most cases; the only statistically important difference was between 1% and 6% of NaOCl.²⁵ In the surface tension study,²⁶ smoother smear layers and less debris were marked when the solution's heat was increased, indicating that heat increased its cleaning capability. The debris particles left on canal walls by 5% NaOCl at 21°C were more intensive than those left by 5% NaOCl at 50°C when used at the same temperature. At 21°C, 2.5% NaOCl produced a similar canal surface appearance to 5% NaOCl, but 2.5% NaOCl at 37°C and 50°C produced smoother smear layers and less debris than 5% NaOCl.²⁷ In,²⁸ 5% NaOCl was evaluated at 21°C and 50°C for debridement capabilities, and it was found that 5% NaOCl heated to 50°C produced a finer particle composition smear layer than was achieved at 21°C. There was a statistically significant difference between results obtained from treatments at different temperatures when teeth were treated with 5.25% NaOCl or 6% NaOCl for 20 min.

2.2. CHX

A CHX solution with a 2% concentration had a lower surface tension and more satisfactory cleansing performance than one with a 0.2% concentration. Because of its cleansing properties, CHX solution should be used as a root canal irrigant at a 2% concentration.²⁹ The surface tension of CHX was lower than that of NaOCl,³⁰

which asserts the results reported in³¹. At³² different temperatures, surface tension in CHX solution at 2% concentration was lower than at 0.2% concentration at both concentrations. Added to a solution, CHX decreases the surface tension of the solution by acting as a cationic active agent. CHX solution with higher concentrations has lower surface tension. Compared with 5% NaOCl, specimens irrigated with 2% CHX solution showed smoother smear layers and fewer debris.³³ A 2% CHX solution at 50°C separated smear layers partially in the coronal and middle thirds of some specimens and left some dentinal tubules with open orifices. In addition, debris-free surfaces have been marked throughout the canal. Despite the similar appearance of canal surfaces between specimens irrigated with 0.2% and 2% CHX solution, 2% CHX produced the best results. There seems to be a correlation between the outcomes and solution surface tensions. 2% CHX solution heated to 50°C cannot entirely remove debris from the apical third.³⁴ This difficulty can be described by the reduction in root canal diameter and subsequent decline in irrigant flow.³⁵ It can be more worthwhile for apical cleaning to use solutions with low surface tension during canal preparations, particularly in narrow canals. In conclusion, both the temperature and concentration of NaOCl and CHX solutions affected surface tension. The surface tension of solutions decreased as the temperature increased; nevertheless, the concentration impact on surface tension was various. At low concentrations of NaOCl, the surface tension decreased, while at low concentrations of CHX, it increased.

2.3. MTAD

A high detergent effect combined with firm antimicrobial effectiveness was the purpose of launching MTAD on the market.³⁶ Study results indicate that MTAD has lower antimicrobial and antifungal activity than NaOCl and CHX in comparison with 5% or 6%^{37,38} The results.³⁹ revealed that MTAD was quite more effective at killing *Escherichia faecalis* than NaOCl when the solutions were diluted.⁴⁰ It is presently used more for final irrigation, after using NaOCl as the main irrigant, than for intra-canal irrigation.⁴¹ A satisfactory wettability along with effective antimicrobial activity is unquestionably crucial for an ideal irrigant solution.⁴² vouched⁴³ that Biopure MTAD has a lower surface tension than NaOCl, even though the NaOCl concentration in this study was half that of an earlier study (5,25%). Low surface tension (34.5 mJ/m²) of MTAD is capable of removing the smear layer and opening the dentinal tubule orifices.⁴⁴ It may therefore be possible to achieve better antibacterial results with these irrigants (having a low surface tension).⁴⁵ Consequently, further research is needed to demonstrate the ability of mentioned fresh irrigants by determining the penetration of antibiotic irrigant solutions into the dentinal tubules, as well as the

decontamination of bacteria with these solutions.⁴⁶

2.4. EDTA

Even though EDTA has been used as an endodontic chelating agent for many years, its antibacterial activities are limited and its surface tension is high.⁴⁷ There have been several ways suggested solving these problems, including adding surfactants and antibiotics (tetracycline) to irrigation solutions. EDTA may be more effective against bacteria when cetrimide (CTR) is added.⁴⁸ It has been reported in⁴⁹ that EDTA compounds with low surface tension alone or in combination with NaOCl can increase root canal dentin's wettability. EDTA solution's surface tension was remarkably reduced after adding surfactant. Root canal dentin had an increase in surface free energy and water contact angle when combined with experimental solutions and NaOCl. When the surfactant is added to the EDTA solution, its surface tension is dramatically declined and its wettability is significantly raised. It is therefore recommended that root canal dentin's adhesion properties may be improved by low-surface tension EDTA solutions. For a prosperous adhesion mechanism, the properties of the adhesive material are also important.⁵⁰

3. Nano-Particles

Despite improving the penetration of NaOCl into the primary canal, surface-active agents do not affect its ability to dissolve pulp tissue. However, there are insufficient data to support a sound conclusion regarding lubrication, antimicrobial, smear layer removal, or debris removal, based on the modification of NaOCl's surface tension. The immediate reaction between the target cells and these particles appears to prevent the organisms from developing resistance^{51,52} Despite their high activity against bacteria, they are compatible with human cells.⁵³ A completely charged imidazolium-based silver nanoparticle (Im AgNPs) showed high antibacterial activity against *Enterococcus faecalis* and cytocompatibility with fibroblasts in a study reported by.⁵⁴ The surface tension of AgNP irrigants is similar to that of distilled water. Compared with the three typical irrigants, the AgNP irrigant has a high surface tension value, which suggests that it might not penetrate dentinal tubules as well. The results of were in agreement with this finding. In this experiment, AgNPs were compared with 2.5% NaOCl and 2% CHX for antimicrobial efficacy against biofilm *E. faecalis* after 5, 15, and 30 minutes under confocal laser scanning microscope conditions. After 5 minutes of irrigation with AgNPs, more viable bacteria were seen in biofilms than in tubules, but after 30 minutes of irrigation, the number of viable bacteria was more prominent in tubules than in biofilms. The scholars finalized that AgNPs could not penetrate dentinal tubules and eliminate bacteria at any time interval tested.

4. Conclusion

Only a few studies have examined the possibility that surface tension may have a bearing on endodontic therapy success. In studies, low-surface tension irrigants have been shown to eliminate the need to clear considerable amounts of root canal dentin to create a debris-free root canal. It would be possible to reduce surface tension so that irrigants could better contact dentinal walls in root canals. As a result, dental irrigants may be adapted more easily to the dentinal walls. NaOCl is more likely to penetrate the primary canal with the help of surface-active agents, according to the evidence available. Both temperature and concentration influenced surface tensions in NaOCl and CHX solutions. The surface tension of both solutions decreased as the temperatures of solutions increased, but the impact of concentration on surface tension was diverse. Low concentrations of NaOCl solution resulted in decreased surface tension, whereas CHX solution resulted in increased surface tension. The im-AgNP solution decreases the surface wettability of dentin and has an evident effect on its physicochemical properties. This solution has a lower viscosity than other irrigants, which allows it to reach root canal apical portions due to its lower viscosity. Because of its higher surface tension, it may not penetrate better inside dentinal tubules. Based on our testing, MTAD appeared to be the best wettability irrigant, followed by 2% CHX and 2.5 percent NaOCl. The wettability of MTAD was the best among the tested irrigants, followed by CHX, 2.5% NaOCl, and two percent of CHX. Surfactants can improve the wetting ability of EDTA and improve root canal dentin adhesion by adding them to EDTA.

5. Source of Funding

None.

6. Conflict of Interest

None.


References

1. Mohammadi Z, Soltani MK, Shalavi S. An update on the management of endodontic biofilms using root canal irrigants and medicaments. *Iran Endod J.* 2014;9(2):89–97.
2. Rosa L, Scolaro GRM, Leanza C, Rapisarda G, Isola S, Cutroneo G. E: Surface tension and wetting ability comparison of sodium hypochlorite and ethylenediaminetetraacetic acid with and without surfactants. *Giornale Italiano Di Endodonzia.* 2021;35(1). doi:10.32067/GIE.2021.35.01.16.
3. Giardino L, Morra M, Becce C. Comparative wettability of different sodium hypochlorite solutions. *G It Endod.* 2012;26(2):57–62.
4. Giardino L, Ambu E, Becce C. Surface tension comparison of four common root canal irrigants and two new irrigants containing antibiotic. *J Endod.* 2006;32(11):1091–4.
5. Almeida LH, Leonardo NG, Gomes AP. Pulp tissue dissolution capacity of sodium hypochlorite combined with cetrimide and polypropylene glycol. *Braz Dent J.* 2013;24(5):477–81.

6. Kwok D, Gietzelt T, Grundke K. Contact angle measurements and contact angle interpretation. Contact angle measurements by axisymmetric drop shape analysis and a goniometer sessile drop technique. *Langmuir*. 1997;13(10):2880–94.
7. Mittal KL. Contact Angle, Wettability and Adhesion. vol. 3. 1st ed. and others, editor. Scrivener Publishing LLC; 2009. p. 520.
8. Cameron JA, Pecora JD, Guimaraes LF, Savioli RN. The effect of a fluorocarbon surfactant on the surface tension of the endodontic irrigant, sodium hypochlorite. A preliminary report. *Aust Dent J*. 1986;31(9):123–30.
9. Rostagni A. Fisica Generale - Meccanica e Termodinamica. Turin, Italy: UTET; 1973. p. 640. Available from: <https://www.goodreads.com/book/show/11307794-fisica-generale---meccanica-e-termodinamica>.
10. Giardino L, Fabbro D, Morra M. Dual Rinse® HEDP increases the surface tension of NaOCl but may increase its dentin disinfection efficacy. *Odontology*. 2019;107(4):521–30.
11. Lottanti S, Gautschi H, Sener B, Zehnder M. Effects of ethylenediaminetetraacetic, etidronic and peracetic acid irrigation on human root dentine and the smear layer. *Int Endod J*. 2009;42(4):335–78.
12. Rath PP, Yiu C, Matinlinna JP, Kishen A, Neelakantan P. The effects of sequential and continuous chelation on dentin. *Dent Mater*. 2020;36(12):1655–65.
13. Lui JN, Kuah HG, Chen NN. Effect of EDTA with and without surfactants or ultrasonics on removal of smear layer. *J Endod*. 2007;33(4):472–7.
14. Taşman F, Cehreli ZC, Oğan C, Etikan I. Surface tension of root canal irrigants. *J Endod*. 2000;26(10):586–93.
15. Neglia R, Ardizzoni A, Giardino L. Comparative in vitro and ex vivo studies on the bactericidal activity of Tetraclean, a new generation endodontic irrigant, and sodium hypochlorite. *New Microbiol*. 2008;31(1):57–65.
16. Mohammadi Z, Mombeinipour A, Giardino L, Shahriari A. Residual antibacterial activity of a new modified sodium hypochlorite-based endodontic irrigation solution. *Med Oral Patol Oral Cir Bucal*. 2011;16(4):588–92.
17. EDTA Plus Brochure, Essential Dental Systems.; 2020. Available from: http://edsdental.com/eds_edta.htm. Accessed on.
18. Estevez R, Conde AJ, De Pablo V. Effect of Passive Ultrasonic Activation on Organic Tissue Dissolution from Simulated Grooves in Root Canals Using Sodium Hypochlorite with or without Surfactants and EDTA. *J Endod*. 2017;43(7):1161–5.
19. Iglesias JE, Pinheiro LS, Weibel DE. Influence of surfactants addition on the properties of calcium hypochlorite solutions. *J Appl Oral Sci*. 2019;27:20180157.
20. Fedele GR, Guastalli AR. Effect of an alcohol-based caries detector on the surface tension of sodium hypochlorite preparations. *Braz Dent J*. 2015;26(1):66–74.
21. Bukiet F, Couderc G, Camps J. Wetting properties and critical micellar concentration of benzalkonium chloride mixed in sodium hypochlorite. *J Endod*. 2012;38(11):1525–34.
22. De-Deus G, Pinho MDB, Reis C. Sodium hypochlorite with reduced surface tension does not improve in situ pulp tissue dissolution. *J Endod*. 2013;39(8):1039–43.
23. Guastalli AR, Clarkson RM, Fedele GR. The effect of surfactants on the stability of sodium hypochlorite preparations. *J Endod*. 2015;41(8):1344–52.
24. Torabinejad M, Handysides R, Khademi A, Bakland LK. Clinical implications of the smear layer in endodontics: a review. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2002;94(6):658–66.
25. Scolaro C, Torrisi L, Cutroneo M. A Liquid Contact Angles on Biocompatible Surfaces in: III^o Workshop Plasmì, Sorgenti, Biofisica ed Applicazioni. 3rd ed. ESE publications, Lecce; 2012. p. 38–42.
26. Stojicic S, Zivkovic S, Qian W. Tissue dissolution by sodium hypochlorite: effect of concentration, temperature, agitation, and surfactant. *J Endod*. 2010;36(9):1558–62.
27. Garbassi F, Morra M, Occhiello E. Polymer surfaces from physics to technology. 1st ed. New York, NY, USA: John Wiley Sons Inc; 1994. p. 510.
28. Yilmaz Z, Basbag B, Buzoglu HD, Gümüsderelioglu M. Effect of low-surface-tension EDTA solutions on the wettability of root canal dentin. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2011;111(1):109–14.
29. Abdollahi-Mansoorkhani HR, Soleimani F, Mahmoudi F. A Multi-Criteria Approach for Comparison of Ginger Extract and Conventional Irrigants in Root Canal Treatment. *Cureus*. 2022;14(9):e29327. doi:10.7759/cureus.29327.
30. Palazzi F, Morra M, Mohammadi Z. Comparison of the surface tension of 5.25% sodium hypochlorite solution with three new sodium hypochlorite-based endodontic irrigants. *Int Endod J*. 2012;45(2):129–64.
31. Fedele GR, Prichard JW, Steier L, De Figueiredo J. The effect of surface tension reduction on the clinical performance of sodium hypochlorite in endodontics. *Int Endod J*. 2013;46(6):492–500.
32. Wang Z, Shen Y, Ma J, Haapasalo M. The effect of detergents on the antibacterial activity of disinfecting solutions in dentin. *J Endod*. 2012;38(7):948–53.
33. Baron A, Lindsey K, Sidow SJ. Effect of a Benzalkonium Chloride Surfactant-Sodium Hypochlorite Combination on Elimination of Enterococcus faecalis. *J Endod*. 2016;42(1):145–54.
34. López P, Conde A, Estevez AJ. Sodium Hypochlorite and a Preparation Containing Glycocholic Acid and Surfactants Have a Synergistic Action on Organic Tissue Dissolution In Vitro. *J Endod*. 2018;44(5):813–8.
35. Palazzi F, Blasi A, Mohammadi Z. Penetration of Sodium Hypochlorite Modified with Surfactants into Root Canal Dentin. *Braz Dent J*. 2016;27(2):208–24.
36. Zehnder M, Schicht O, Sener B, Schmidlin P. Reducing surface tension in endodontic chelator solutions has no effect on their ability to remove calcium from instrumented root canals. *J Endod*. 2005;31(8):590–2.
37. Farshad M, Abbaszadegan A, Ghahramani Y, Jamshidzadeh A. Effect of Imidazolium-Based Silver Nanoparticles on Root Dentin Roughness in Comparison with Three Common Root Canal Irrigants. *Iran Endod J*. 2017;12(1):83–6.
38. Wu D, Fan W, Kishen A, Gutmann JL, Fan B. Evaluation of the antibacterial efficacy of silver nanoparticles against Enterococcus faecalis biofilm. *J Endod*. 2014;40(2):285–90.
39. Bruniera JFB, Silva-Sousa YTC, Lara MG, Pitondo-Silva A, Marcaccini AM, Miranda CES. Development of Intracanal Formulation Containing Silver Nanoparticles. *Braz Dent J*. 2014;25(4):302–8.
40. Abbaszadegan A, Nabavizadeh M, Gholami A, Aleyasin Z, Dorostkar S, Saliminasab M. Positively charged imidazolium-based ionic liquid-protected silver nanoparticles: a promising disinfectant in root canal treatment. *Int Endod J*. 2015;48(8):790–800.
41. Abbaszadegan A, Ghahramani Y, Gholami A, Hemmateenejad B, Dorostkar S, Nabavizadeh M. The effect of charge at the surface of silver nanoparticles on antimicrobial activity against gram-positive and gram-negative bacteria: a preliminary study. *J Nanomater*. 2015;2015(720654):53.
42. Rai M, Deshmukh S, Ingle A, Gade A. Silver nanoparticles: the powerful nanoweapon against multidrug-resistant bacteria. *J Appl Microbiol*. 2012;112(5):841–52.
43. Abbaszadegan A, Gholami A, Abbaszadegan S, Aleyasin ZS, Ghahramani Y, Dorostkar S, et al. The Effects of Different Ionic Liquid Coatings and the Length of Alkyl Chain on Antimicrobial and Cytotoxic Properties of Silver Nanoparticles. *Iran Endod J*. 2017;12(4):481–7.
44. Ghahramani Y, Yaghoubi F, Motamedi R, Jamshidzade A, Abbaszadegan A. Effect of Endodontic Irrigants and Medicaments Mixed with Silver Nanoparticles against Biofilm Formation of Enterococcus faecalis. *Iran Endod J*. 2018;13(4):559–64.
45. Slawson RM, Trevors JT, Lee H. Silver accumulation and resistance in *Pseudomonas stutzeri*. *Arch Microbiol*. 1992;158(6):398–404.
46. Zhao G. Multiple parameters for the comprehensive evaluation of the susceptibility of *Escherichia coli* to the silver ion. *Biometals*.

- 1998;11(1):27–32.
47. Prabhu S, Poulouse EK. Silver nanoparticles: mechanism of antimicrobial action, synthesis, medical applications, and toxicity effects. *Int Nano Lett.* 2012;2(1):1–10.
48. Nabavizadeh M, Ghahramani Y, Abbaszadegan A, Jamshidzadeh A, Jenabi P, Makarempour A. In Vivo Biocompatibility of an Ionic Liquid-protected Silver Nanoparticle Solution as Root Canal Irrigant. *Iran Endod J.* 2018;13(3):293–8.
49. Machnick TK, Munoz C, Shabahang S, Torabinejad M. Effect of MTAD on flexural strength and modulus of elasticity of dentin. *J Endod.* 2003;29(11):747–50.
50. Torabinejad M, Cho Y, Khademi AA, Bakland LK, Shabahang S. The effect of various concentrations of sodium hypochlorite on the ability of MTAD to remove the smear layer. *J Endod.* 2003;29(11):233–42.
51. Machnick TK, Torabinejad M, Munoz CA, Shabahang S. Effect of MTAD on the bond strength to enamel and dentin. *J Endod.* 2003;29(11):818–39.
52. Sumbul M, Çolak KM, Arslan H. Effect of different temperature and pH values of EDTA on sealer penetration into dentinal tubules: a pilot study with confocal laser scanning microscope. *J Dent Fac Atatürk Uni.* 2021;31(3):379–84.
53. Aslantas EE, Buzoglu HD, Altundasar E, Serper A. Effect of EDTA, sodium hypochlorite, and chlorhexidine gluconate with or without surface modifiers on dentin microhardness. *J Endod.* 2014;40(6):876–85.
54. Gopikrishna PV, Kumar AP, Narayanan LL. Influence of temperature and concentration on the dynamic viscosity of sodium hypochlorite in comparison with 17% EDTA and 2% chlorhexidine gluconate: An in vitro study. *J Conserv Dent.* 2014;17(1):57–60.

Author biography

Hamidreza Abdollahi Mansoorkhani, DDS  <https://orcid.org/0000-0003-4107-7891>

Farshid Mahmoudi, Assistant Professor

Cite this article: Mansoorkhani HA, Mahmoudi F. The importance of surface tension in endodontic irrigation: A review study. *Arch Dent Res* 2022;12(2):76-80.